

**In the Claims**

Applicants have submitted a new complete claim set showing marked-up claims with insertions indicated by underlining and deletions indicated by strikeouts and/or double bracketing.

Please amend claims 12, 15, 18-21, 23-33, 36-39, 56, 77, and 84 as indicated below:

---

1. (Original): A method for desymmetrization, comprising:

Bl providing a catalyst and a molecular substrate having a plane of symmetry, the catalyst being present in an amount of less than 15 mol%, relative to an amount of substrate; and causing an olefin metathesis reaction involving the molecular substrate to occur to form a product free of a plane of symmetry.

2. (Cancelled)

3. (Original): A method as in claim 1, wherein the molecular substrate is selected from the group consisting of achiral and meso substrates.

4. (Original): A method as in claim 1, wherein the molecular substrate is selected from the group consisting of cyclic and acyclic substrates.

5. (Original): A method as in claim 1, wherein the product is selected from the group consisting of cyclic and acyclic products.

6. (Original): A method as in claim 1, wherein the product includes at least one ring having a ring size of less than about 20 atoms.


7. (Original): A method as in claim 1, wherein the product includes at least one ring having a ring size of less than about 10 atoms.

8. (Cancelled)

10. (Original): A method as in claim 1, wherein the catalyst is present in an amount of less than about 5 % mol.

11. (Original): A method as in claim 1, wherein the catalyst is present in an amount of less than about 1 mol%.

12. (Currently Amended): A method as in claim 1, wherein the molecular substrate comprises a structure:




wherein "1" and "2" can be the same or different and each of "1" and "2" denotes a bond selected from the group consisting of a double bond and a triple bond; a, b, d, and e can be the same or different and each of a, b, d and e is an integer equaling 0 to 1; m and n can be the same or different and each of m and n are integers equaling 0-20; A, B, D, E and R<sup>1</sup> - R<sup>3</sup> can be the same or different and each of A, B, D, E and R<sup>1</sup> - R<sup>3</sup> is selected from the group consisting of hydrogen, hydroxy, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl, wherein C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

13. (Original): A method as in claim 12, wherein each of m and n are integers equaling 0-10.

14. (Original): A method as in claim 12, wherein the functional group including at least one non-carbon element is selected from the group consisting of O, S, Se, silane, silyl ether, carbonyl, carboxyl, carboxylate, ether, ester, anhydride, acyl, cyano, NO<sub>2</sub>, alkyloxy, aryloxy, hydroxy, hydroxyalkyl, amino, alkylamino, arylamino, amido, thioalkyl, thioaryl, sulfonate, phosphate, phosphonate, phosphane and stannane.

15. (Original): A method as in claim 1, wherein the molecular substrate comprises a structure:




wherein "1" and "2" can be the same or different and each of "1" and "2" denotes a bond selected from the group consisting of a double bond and a triple bond; a, b, d, and e can be the same or different and each of a, b, d and e is an integer equaling 0 to 1; m and n can be the same or different and each of m and n are integers equaling 0-20; A, B, D, E and R<sup>1</sup> - R<sup>3</sup> can be the same or different and each of A, B, D, E and R<sup>1</sup> - R<sup>3</sup> is selected from the group consisting of hydrogen, hydroxy, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>3</sub>-C<sub>20</sub> aryl and C<sub>2</sub>-C<sub>20</sub> alkynyl, wherein C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>3</sub>-C<sub>20</sub> aryl and C<sub>2</sub>-C<sub>20</sub> alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

16. (Original): A method as in claim 15, wherein each of m and n are integers equaling 0-10.

17. (Original): A method as in claim 15, wherein the functional group including at least one non-carbon element is selected from the group consisting of O, S, Se, silane, silyl ether, carbonyl, carboxyl, carboxylate, ether, ester, anhydride, acyl, cyano, NO<sub>2</sub>, alkyloxy, aryloxy, hydroxy, hydroxyalkyl, amino, alkylamino, arylamino, amido, thioalkyl, thioaryl, sulfonate, phosphate, phosphonate, phosphane and stannane.

bl 18. (Currently Amended): A method as in claim 15, wherein X is selected from the group consisting of  $\text{CR}^8\text{R}^9$ , carbonyl, ester,  $\text{SiR}^8\text{R}^9$ ,  $\text{OSi}(\text{R}^8)(\text{R}^9)$ ,  $\text{SnR}^8\text{R}^9$ , O, S, Se,  $\text{NR}^8$ ,  $\text{PR}^8$  and  $\text{PO}_3\text{R}^8$ ;  $\text{R}^8$  and  $\text{R}^9$  can be the same or different and each of  $\text{R}^8$  and  $\text{R}^9$  is selected from the group consisting of hydrogen,  $\text{C}_1\text{-C}_{20}$  alkyl,  $\text{C}_{42}\text{-C}_{20}$  alkenyl,  $\text{C}_{43}\text{-C}_{20}$  aryl and  $\text{C}_{42}\text{-C}_{20}$  alkynyl, wherein  $\text{C}_1\text{-C}_{20}$  alkyl,  $\text{C}_{42}\text{-C}_{20}$  alkenyl,  $\text{C}_{43}\text{-C}_{20}$  aryl and  $\text{C}_{42}\text{-C}_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.


19. (Currently Amended): A method as in claim 15, wherein the molecular substrate comprises a structure:



wherein  $\text{R}^4 - \text{R}^7$  can be the same or different and each of  $\text{R}^4 - \text{R}^7$  is selected from the group consisting of hydrogen, hydroxy,  $\text{C}_1\text{-C}_{20}$  alkyl,  $\text{C}_{42}\text{-C}_{20}$  alkenyl,  $\text{C}_{43}\text{-C}_{20}$  aryl and  $\text{C}_{42}\text{-C}_{20}$  alkynyl, wherein  $\text{C}_1\text{-C}_{20}$  alkyl,  $\text{C}_{42}\text{-C}_{20}$  alkenyl,  $\text{C}_{43}\text{-C}_{20}$  aryl and  $\text{C}_{42}\text{-C}_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

20. (Currently Amended): A method as in claim 19, wherein X is selected from the group consisting of  $\text{CR}^8\text{R}^9$ , carbonyl, ester,  $\text{SiR}^8\text{R}^9$ ,  $\text{SiR}^8\text{R}^9$ ,  $\text{OSi}(\text{R}^8)(\text{R}^9)$ ,  $\text{SnR}^8\text{R}^9$ , O, S, Se,  $\text{NR}^8$ ,  $\text{PR}^8$ , and  $\text{PO}_3\text{R}^8$ ;  $\text{R}^8$  and  $\text{R}^9$  can be the same or different and each of  $\text{R}^8$  and  $\text{R}^9$  is selected from the group consisting of hydrogen,  $\text{C}_1\text{-C}_{20}$  alkyl,  $\text{C}_{42}\text{-C}_{20}$  alkenyl,  $\text{C}_{43}\text{-C}_{20}$  aryl and  $\text{C}_{42}\text{-C}_{20}$  alkynyl, wherein  $\text{C}_1\text{-C}_{20}$  alkyl,  $\text{C}_{42}\text{-C}_{20}$  alkenyl,  $\text{C}_{43}\text{-C}_{20}$  aryl and  $\text{C}_{42}\text{-C}_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

21. (Currently Amended): A method as in claim 1, wherein the molecular substrate comprises a structure:




wherein "1" and "2" can be the same or different and each of "1" and "2" denotes a bond selected from the group consisting of a double bond and a triple bond; X and Y can be the same or different and each is a functional substituent; a, b, d, e and g can be the same or different and each of a, b, d, e and g are integers equaling 0 to 1; m, n, o and p can be the same or different and each of m, n, o and p are integers equaling 0-20; A, B, D, E, G and  $R^1 - R^3$  can be the same or different and each of A, B, D, E, G and  $R^1 - R^3$  is selected from the group consisting of hydrogen, hydroxy,  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl, and wherein  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl, wherein and  $C_{42}$ - $C_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

22. (Original): A method as in claim 21, wherein each of m and n are integers equaling 0-10.

23. (Currently Amended): A method as in claim 21, wherein X and Y are selected from the group consisting of  $CR^9R^{10}$ , carbonyl, ester,  $SiR^9R^{10}$ ,  $OSi(R^9)(R^{10})$ ,  $SnR^9R^{10}$ , B, O, S, Se,  $NR^9$ ,  $PR^9$  and  $PO_3R^9$ ;  $R^9$  and  $R^{10}$  can be the same or different and each of  $R^9$  and  $R^{10}$  is selected from the group consisting of hydrogen,  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl, wherein  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.


24. (Currently Amended): A method as in claim 21, wherein the molecular substrate comprises a structure:



wherein  $R^4 - R^8$  can be the same or different and each of  $R^4 - R^8$  is selected from the group consisting of hydrogen, hydroxy,  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl, wherein  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

25. (Currently Amended): A method as in claim 24, wherein X is selected from the group consisting of  $CR^9R^{10}$ , carbonyl, ester,  $SiR^9R^{10}$ ,  $OSi(R^9)(R^{10})$ ,  $SnR^9R^{10}$ , B, O, S, Se,  $NR^9$ ,  $PR^9$  and  $PO_3R^9$ ;  $R^9$  and  $R^{10}$  can be the same or different and each of  $R^9$  and  $R^{10}$  is selected from the group consisting of hydrogen,  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl, wherein  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.


26. (Currently Amended): A method as in claim 1, wherein the molecular substrate comprises a structure:



wherein "1" and "2" can be the same or different and each of "1" and "2" denotes a bond selected from the group consisting of a double bond and a triple bond; X and Y can be the same or different and each is a functional substituent; a, b, d and e can be the same or different and each of a, b, d and e are integers equaling 0 to 1; m, n, o, p and q can be the same or different and each of m, n, o, p and q are integers equaling 0-20; A, B, D, E and  $R^1 - R^4$  can be the same or different and each of A, B, D, E and  $R^1 - R^4$  is selected from the group consisting of hydrogen, hydroxy,  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl, wherein  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

27. (Currently Amended): A method as in claim 26, wherein X and Y are selected from the group consisting of  $CR^9R^{10}$ , carbonyl, ester,  $SiR^9R^{10}$ ,  $OSi(R^9)(R^{10})$ ,  $SnR^9R^{10}$ , B, O, S, Se,  $NR^9$ ,  $PR^9$  and  $PO_3R^9$ ;  $R^9$  and  $R^{10}$  can be the same or different and each of  $R^9$  and  $R^{10}$  is selected from the group consisting of hydrogen,  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl, wherein  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

28. (Currently Amended): A method as in claim 26, wherein the molecular substrate comprises a structure:




wherein  $R^5 - R^8$  can be the same or different and each of  $R^5 - R^8$  is a selected from the group consisting of hydrogen, hydroxy,  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl, wherein  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl, wherein and  $C_{42}$ - $C_{20}$  alkynyl are hydrocarbons optionally interrupted a functional group including at least one non-carbon element.

29. (Currently Amended): A method as in claim 28, wherein X is selected from the group consisting of  $CR^9R^{10}$ , carbonyl, ester,  $SiR^9R^{10}$ ,  $OSi(R^9)(R^{10})$ ,  $SnR^9R^{10}$ , B, O, S, Se,  $NR^9$ ,  $PR^9$  and  $PO_3R^9$ ;  $R^9$  and  $R^{10}$  can be the same or different and each of  $R^9$  and  $R^{10}$  is selected from the group consisting of hydrogen,  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl, wherein  $C_1$ - $C_{20}$  alkyl,  $C_{42}$ - $C_{20}$  alkenyl,  $C_{43}$ - $C_{20}$  aryl and  $C_{42}$ - $C_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

30. (Currently Amended): A method as in claim 1, wherein the molecular substrate comprises a structure:






wherein "■" denotes a bond selected from the group consisting of a double bond and a triple bond; V, W, X, Y and Z can be the same or different and V, W, X, Y and Z are functional substituents; a and b can be the same or different and each of a and b are integers equaling 0 to 1; m, n, o, p, q and r can be the same or different and each of m, n, o, p, q and r are integers equaling 0-20; A, B and  $R^1 - R^3$  can be the same or different and each of A, B and  $R^1 - R^3$  is selected from the group consisting of hydrogen, hydroxy,  $C_1-C_{20}$  alkyl,  $C_{42}-C_{20}$  alkenyl,  $C_{43}-C_{20}$  aryl and  $C_{42}-C_{20}$  alkynyl, wherein  $C_1-C_{20}$  alkyl,  $C_{42}-C_{20}$  alkenyl,  $C_{43}-C_{20}$  aryl and  $C_{42}-C_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

31. (Currently Amended): A method as in claim 30, wherein each of V, W, X, Y and Z is selected from the group consisting of  $CR^6R^7$ , carbonyl, ester,  $SiR^6R^7$ ,  $OSi(R^6)(R^7)$ ,  $SnR^6R^7$ , B, O, S, Se,  $NR^6$ ,  $PR^6$  and  $PO_3R^6$ ;  $R^6$  and  $R^7$  can be the same or different and each of  $R^6$  and  $R^7$  is selected from the group consisting of hydrogen,  $C_1-C_{20}$  alkyl,  $C_{42}-C_{20}$  alkenyl,  $C_{43}-C_{20}$  aryl and  $C_{42}-C_{20}$  alkynyl, wherein  $C_1-C_{20}$  alkyl,  $C_{42}-C_{20}$  alkenyl,  $C_{43}-C_{20}$  aryl and  $C_{42}-C_{20}$  alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

32. (Currently Amended): A method as in claim 30, wherein the molecular substrate comprises a structure:




wherein R<sup>4</sup> and R<sup>5</sup> can be the same or different and each of R<sup>4</sup> and R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl, wherein C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

33. (Currently Amended): A method as in claim 32, wherein each of X, Y and Z is selected from the group consisting of CR<sup>6</sup>R<sup>7</sup>, carbonyl, ester, SiR<sup>6</sup>R<sup>7</sup>, OSi(R<sup>6</sup>)(R<sup>7</sup>), SnR<sup>6</sup>R<sup>7</sup>, B, O, S, Se, NR<sup>6</sup>, PR<sup>6</sup> and PO<sub>3</sub>R<sup>6</sup>; R<sup>6</sup> and R<sup>7</sup> can be the same or different and each of R<sup>6</sup> and R<sup>7</sup> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl, wherein C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.


34. (Original): A method as in claim 1, wherein the olefin metathesis reaction is selected from the group consisting of ring-closing metathesis and ring-opening metathesis.

35. (Original): A method as in claim 1, wherein the molecular substrate is a first molecular substrate, the method further comprising a second molecular substrate and the olefin metathesis reaction is a cross-metathesis reaction.

36. (Currently Amended): A method as in claim 35, wherein the first molecular substrate is selected from the group consisting of:



and the second molecular substrate comprises a structure:




wherein "1" and "2" can be the same or different and each of "1" and "2" denotes a bond selected from the group consisting of a double bond and a triple bond; W and X can be the same or different and W and X are functional substituents; a, b, d and e can be the same or different and each of a, b, d and e are integers equaling 0 to 1; m, n, o, p, q and r can be the same or different and each of m, n, o, p, q and r are integers equaling 0-20; A, B, D, E and R<sup>1</sup> - R<sup>3</sup> can be the same or different and each of A, B, D, E and R<sup>1</sup> - R<sup>3</sup> is selected from the group consisting of hydrogen, hydroxy, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl, wherein C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element; Y and Z can be the same or different and each of Y and Z is selected from the group consisting of CN, carboxylic ester, amide, acid, halogen, hydrogen, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl, wherein C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl are hydrocarbons optionally interrupted a functional group including at least one non-carbon element.


37. (Currently Amended): A method as in claim 36, wherein each of W and X is selected from the group consisting of CR<sup>8</sup>R<sup>9</sup>, carbonyl, ester, SiR<sup>8</sup>R<sup>9</sup>, OSi(R<sup>8</sup>)(R<sup>9</sup>), SnR<sup>8</sup>R<sup>9</sup>, O, S, Se, NR<sup>8</sup>, PR<sup>8</sup> and PO<sub>3</sub>R<sup>8</sup>; R<sup>8</sup> and R<sup>9</sup> can be the same or different and each of R<sup>8</sup> and R<sup>9</sup> is selected

B1  
from the group consisting of hydrogen, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl, wherein C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

38. (Currently Amended): A method as in claim 36, wherein the first molecular substrate is selected from the group consisting of:



and the second molecular substrate comprises a structure:



wherein R<sup>4</sup> - R<sup>7</sup> can be the same or different and each of R<sup>4</sup> - R<sup>7</sup> is selected from the group consisting of hydrogen, hydroxy, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl, wherein C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

39. (Currently Amended): A method as in claim 38, wherein each of W and X is selected from the group consisting of CR<sup>8</sup>R<sup>9</sup>, carbonyl, ester, SiR<sup>8</sup>R<sup>9</sup>, OSi(R<sup>8</sup>)(R<sup>9</sup>), SnR<sup>8</sup>R<sup>9</sup>, O, S, Se, NR<sup>8</sup>, PR<sup>8</sup> and PO<sub>3</sub>R<sup>8</sup>; R<sup>8</sup> and R<sup>9</sup> can be the same or different and each of R<sup>8</sup> and R<sup>9</sup> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl, wherein C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>42</sub>-C<sub>20</sub> alkenyl, C<sub>43</sub>-C<sub>20</sub> aryl and C<sub>42</sub>-C<sub>20</sub> alkynyl are hydrocarbons optionally interrupted by a functional group including at least one non-carbon element.

40. (Original): A method as in claim 1, wherein the product is formed at a turnover number of at least about 5, the product being at least one enantiomer formed in an enantiomeric excess of at least about 20%.

41. (Original): A method as in claim 40, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 50%.

42. (Original): A method as in claim 40, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 85%.

43. (Original): A method as in claim 40, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 90%.

44. (Original): A method as in claim 40, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 95%.

45. (Original): A method as in claim 40, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 99%.

46. (Original): A method as in claim 40, wherein two enantiomers are each formed in an enantiomeric excess of at least about 20%.

47. (Original): A method as in claim 46, wherein the two enantiomers are each formed in an enantiomeric excess of at least about 50%.

48. (Original): A method as in claim 46, wherein the two enantiomers are each formed in an enantiomeric excess of at least about 85%.

49. (Original): A method as in claim 46, wherein the two enantiomers are each formed in an enantiomeric excess of at least about 90%.

50. (Original): A method as in claim 46, wherein the two enantiomers are each formed in an enantiomeric excess of at least about 95%.

B1  
51. (Original): A method as in claim 46, wherein the two enantiomers are each formed in an enantiomeric excess of at least about 99%.

52. (Original): A method as in claim 40, wherein the turnover number is at least about 10.

53. (Original): A method as in claim 40, wherein the turnover number is at least about 25.

54. (Original): A method as in claim 40, wherein the turnover number is at least about 50.


55. (Original): A method as in claim 40, wherein the turnover number is at least about 100.

56.(Currently Amended): A method as in claim 2 1, wherein the catalyst is a metal complex.

57. (Original): A method as in claim 56, wherein the metal complex is a transition metal complex including at least one metal-carbon double bond.

58. (Original): A method as in claim 57, wherein the metal complex is a transition metal dialkoxide complex.

59. (Original): A method as in claim 58, wherein the dialkoxide complex comprises a structure:



wherein the catalyst has a chiral dialkoxide ligand, denoted by  $\begin{pmatrix} \text{O} \\ \text{O} \end{pmatrix}$ , the dialkoxide being of at least 80 % optical purity, M is a transition metal ion, and  $R^1$  and  $R^2$  can be the same or different, and each is selected from the group consisting of  $C_1$ - $C_{12}$  alkyl, heteroalkyl, aryl, heteroaryl and adamantyl.

60. (Original): A method as in claim 59, wherein  $R^1$  is selected from the group consisting of 2,6-dimethylphenyl, 2,6-diethylphenyl and 2,6-diisopropylphenyl and  $R^2$  is selected from the group consisting of methyl, ethyl and phenyl.

61. (Original): A method for desymmetrization, comprising:  
providing a catalyst and a molecular substrate having a plane of symmetry; and  
allowing an olefin metathesis desymmetrization reaction to occur in the absence of solvent to form a product free of a plane of symmetry.

62. (Original): A method as in claim 61, wherein the catalyst is present in an amount of less than 15 mol%, relative to an amount of substrate.

63. (Original): A method as in claim 61, wherein the catalyst is present in an amount of less than 10 mol%, relative to an amount of substrate.

64. (Original): A method as in claim 61, wherein the catalyst is present in an amount of less than 5 mol%, relative to an amount of substrate.

65. (Original): A method as in claim 61, wherein the olefin metathesis reaction is selected from the group consisting of a ring-closing and a ring-opening reaction.

66. (Original): A method as in claim 61, wherein the molecular substrate is a first molecular substrate, the method further comprising a second molecular substrate and the olefin metathesis reaction is a cross-metathesis reaction.

67. (Original): A method as in claim 61, wherein the desymmetrization reaction causes at least one enantiomer of a product to form in an enantiomeric excess of at least about 20% at a turnover number of at least about 5.

68. (Original): A method as in claim 67, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 50%.

69. (Original): A method as in claim 67, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 85%.

70. (Original): A method as in claim 67, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 90%.

71. (Original): A method as in claim 67, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 95%.

72. (Original): A method as in claim 67, wherein the at least one enantiomer is formed in an enantiomeric excess of at least about 99%.

73. (Original): A method as in claim 67, wherein two enantiomers are formed in an enantiomeric excess of at least about 20%.


74. (Original): A method for catalytic desymmetrization, comprising:  
providing a molecular substrate having a plane of symmetry and a catalyst, the catalyst being present in an amount of less than 15 mol%, relative to an amount of substrate; and  
allowing a desymmetrization reaction to occur to form a product having a quaternary carbon center in at least about 20% enantiomeric excess.


75. (Original): A method as in claim 74, wherein the desymmetrization reaction is a carbon-carbon bond forming reaction.





76. (Original): A method as in claim 75, wherein the desymmetrization reaction is an olefin metathesis reaction.

β1 77. (Currently Amended): A composition comprising a structure:



wherein M is a metal ion and  is a chiral dialkoxide of at least 80 % optical purity, the

dialkoxide having sufficient rigidity such that a  reaction site is of sufficient shape specificity, defined in part by the dialkoxide and a M=N-R site, to cause a molecular substrate

having a plane of symmetry to react with a M=C center at the  reaction site,

forming a catalytic olefin metathesis product that has at least a 50 % enantiomeric excess of at least one enantiomer present in the mixture, the product being free of a plane of symmetry,

wherein R<sup>2</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>12</sub> alkyl, heteroalkyl, aryl, heteroaryl and adamantyl.

78. (Original): A method for performing a kinetic resolution, comprising:

providing at least one substrate having at least one olefin group, the substrate having a plane of symmetry;

selecting a catalyst of sufficient steric bulk to initiate an olefin metathesis desymmetrization reaction involving the at least one substrate to achieve a  $k_{rel}$  of at least about 10.

79. (Original): A method as in claim 78, wherein the reaction is selected from the group consisting of a ring-opening metathesis reaction, a cross-metathesis reaction and a ring-closing metathesis reaction.


b1  
80. (Original): A method for performing an asymmetric olefin metathesis reaction, comprising:  
providing a substrate comprising at least one olefin group associated with a ring structure,  
the substrate having a plane of symmetry;  
reacting a catalyst with the substrate to initiate an olefin metathesis desymmetrization  
reaction to achieve a  $k_{rel}$  of at least about 5.

81. (Original): A method as in claim 80, wherein the reaction further comprises a kinetic  
resolution.


82. (Original): A method for performing an asymmetric olefin metathesis reaction, comprising:  
providing two substrates, at least one substrate having a plane of symmetry and each  
substrate containing at least one olefin group;  
reacting a catalyst with the substrates to form a product free of a plane of symmetry  
having an enantiomeric excess of at least about 50%.

83. (Original): A method as in claim 82, wherein the reaction is selected from the group  
consisting of a ring-opening metathesis reaction, a cross-metathesis reaction, kinetic resolution  
and a combination thereof.

84. (Currently Amended): A method as in any one of claims 78, 80 or 82 wherein the catalyst  
comprises a structure:




wherein the catalyst has a chiral dialkoxide ligand, denoted by



the alkoxide being of at least 80% optical purity, M is a transition metal ion, and R<sup>1</sup> and each R<sup>2</sup>

can be the same or different, and each is selected from the group consisting of C<sub>1</sub>-C<sub>12</sub> alkyl, heteroalkyl, aryl, heteroaryl and adamantyl.

85. (Original): A method as in claim 84, wherein the catalyst comprises a structure:




wherein R<sup>1</sup> - R<sup>3</sup> can be the same or different and each is selected from the group consisting of hydrogen, alkyls, aryls, alkaryl and substituted derivatives thereof.

86. (Original): A method as in claim 85, wherein R<sup>3</sup> is selected from the group consisting of ethyl, *i*-Pr, *t*-Bu and adamantyl and R<sup>1</sup> and R<sup>2</sup> selected from the group consisting of *i*-Pr and methyl.

87. (Original): A method as in claim 85, wherein R<sup>1</sup> is CF<sub>3</sub> and R<sup>2</sup> is hydrogen.

88. (Original): A method as in claim 84, wherein the catalyst comprises a structure:



wherein R<sup>1</sup> - R<sup>4</sup> can be the same or different and each is selected from the group consisting of hydrogen, alkyls, aryls, alkaryl and substituted derivatives thereof.

89. (Original): A method as in claim 88, wherein  $R^3$  is selected from the group consisting of 2,4,6-tri(*i*-propyl)phenyl, phenyl and *t*-Bu,  $R^1$  and  $R^2$  are selected from the group consisting of *i*-Pr and methyl and  $R^4$  is selected from the group consisting of hydrogen and *t*-Bu.

90. (Original): A method as in claim 88, wherein  $R^1$  is  $CF_3$  and  $R^2$  is hydrogen.

